WO 2005/054444 PCT/KR2004/003024

## What is claimed is:

- 1. A chemically defined medium for fermentation culture of a strain of the genus Candida, which comprises 1-10 g/ l of urea, 1-10 g/ l of potassium diphosphate, 0.01-1 g/ l of magnesium sulfate, 0.1-10 mg/ l of MnSO<sub>4</sub> · 4H<sub>2</sub>O, 0.1-10 mg/ l of CoCl<sub>2</sub> · 6H<sub>2</sub>O, 0.1-10 mg/ l of NaMoO<sub>4</sub> · 2H<sub>2</sub>O, 0.1-10 mg/ l of ZnSO<sub>4</sub> · 7H<sub>2</sub>O, 0.1-10 mg/ l of AlCl<sub>3</sub> · 6H<sub>2</sub>O, 0.1-10 mg/ l of CuCl<sub>2</sub> · 2H<sub>2</sub>O, 0.01-5 mg/ l of H<sub>3</sub>BO<sub>3</sub>, 1-100 mg/ l of FeSO<sub>4</sub> · 7H<sub>2</sub>O, 0.1-10 mg/ l of ascorbic acid, 1-100 mg/ l of biotin, 1-100 mg/ l of choline, and 0.1-10 mg/ l of pyridoxine.
- 2. A process for producing xylitol in high yield by recycling culture of a strain of the genus *Candida*, which comprises the steps of:

inoculating the strain in a xylose-containing medium and culturing the strain in the xylose-containing medium in a bioreactor;

releasing a culture from the bioreactor and introducing a fresh xylose15 containing medium to the bioreactor continuously;

separating the strain and a culture filtrate from the culture; and recycling the strain to the bioreactor and recovering xylitol from the culture filtrate.

- 20 3. The process of claim 2, wherein the strain of the genus *Candida* is *Candida tropicalis* or its mutant strain.
  - 4. The process of claim 2, wherein the xylose-containing medium is the chemically defined medium of claim 1 or a complex medium.

WO 2005/054444 PCT/KR2004/003024

5. The process of claim 2, wherein the culturing is performed by a fedbatch culture or a batch culture.

- 6. The process of claim 5, wherein in the fed-batch culture, the medium is
  5 gradually supplemented with xylose so that the concentration of xylose is maintained at
  40-50g/ \ell\$ on the basis of the medium.
  - 7. The process of any one of claims 2, 4, 5, and 6, wherein the culturing is performed at an agitation speed of 400-600rpm.

8. The process of claim 2, wherein the separation of the strain and the culture filtrate from the culture is performed by a vacuum microfiltration system or a centrifuge.

10

9. The process of claim 2 or 8, wherein the separated strain is concentrated to a density of 10-100g/l and recycled.